

Semigroup Forum (2012) 85:513–524
 DOI 10.1007/s00233-012-9430-2

RESEARCH ARTICLE

Automorphisms of partition order-decreasing transformation monoids

Haobo Yang · Xiuliang Yang

Received: 24 January 2010 / Accepted: 18 August 2012 / Published online: 13 September 2012
 © The Author(s) 2012. This article is published with open access at Springerlink.com

Abstract Let T_n be the full transformation semigroup on a finite set $X_n = \{1, 2, \dots, n\}$. Let ρ be an equivalence relation on X_n and \preceq be a total order on the partition set X_n/ρ . We describe all automorphisms of the partition order-decreasing transformation monoid:

$$T(\rho, \preceq) = \{\alpha \in T_n : (x\alpha)\rho \preceq x\rho, \forall x \in X_n\}$$

that generalizes the results of Schreier (Fundam. Math., 28:261–264, 1936) and Šutov (Izv. Vysš. Učebn. Zaved., Mat., 3:177–184, 1961).

Keywords Transformation semigroup · Partition order-decreasing · Automorphism group

1 Introduction

For the standard definitions on semigroups and transformation semigroups we refer the reader to the books [3–5].

Let $X_n = \{1, 2, \dots, n\}$. A submonoid M of the semigroup T_n of full transformations on X_n is *intransitive* if there exist x, y in X_n such that $(x)\varphi \neq y$ for any $\varphi \in M$.

Communicated by Jean-Eric Pin.

This work is supported by NSF (China) grant no. 10971086.

H. Yang
 Qiangjiang College, Hangzhou Normal University, Hangzhou City, Zhejiang 310012, P.R. China
 e-mail: yanghb@hznu.edu.cn

X. Yang (✉)
 Department of Mathematics, Hangzhou Normal University, Xiasha, Hangzhou City, Zhejiang 310036, P.R. China
 e-mail: yxl@hznu.edu.cn

A submonoid M of T_n said to be *half-transitive* provided that it is intransitive, and for every ordered pair $(x, y) \in X_n \times X_n$ there is some $\varphi \in M$ such that either $x\varphi = y$ or $y\varphi = x$. In [10] we showed that, for every half-transitive submonoid M of T_n , there exist a non-universal equivalence relation ρ on X_n and a total order \preceq on the partition set X_n/ρ such that M lies inside a half-transitive submonoid $T(\rho, \preceq)$ of T_n defined by

$$T(\rho, \preceq) = \{\alpha \in T_n : (x\alpha)\rho \preceq x\rho, \forall x \in X_n\}.$$

Here we consider the monoid $T(\rho, \preceq)$ for arbitrary equivalence relation ρ on X_n . In particular, if ρ is universal then $T(\rho, \preceq) = T_n$; $T(\rho, \preceq)$ is the order-decreasing finite full transformation monoid if ρ is the identity relation [9]. If $X_n/\rho = \{\{1\}, X_n - \{1\}\}$ and $\{1\} \prec X_n - \{1\}$ then $T(\rho, \preceq)$ is isomorphic to PT_{n-1} , the semigroup of partial transformations of $X_n - \{1\}$.

Note that automorphisms of T_n and PT_{n-1} were described by Schreier and Šutov in [6] and [8] respectively. In present paper, we will describe automorphisms of the monoid $T(\rho, \preceq)$. Thus our result generalizes the results of Schreier [6] and Šutov [8].

For the topics of automorphisms of transformation semigroups we refer the reader to the references of [1] and [3]. We point out that $T(\rho, \preceq)$ is not the centralizer of any idempotent in T_n [1] and, in general, $T(\rho, \preceq)$ does not contain all constant transformations of X_n . Our method is different from any one of the references of [1] and [7]. Here we use a basic fact that an automorphism φ of a monoid S maps units of S to its units, and idempotents of S to its idempotents.

To achieve our aims, the organization of the paper is as follows: In Sect. 2 we investigate the units and the idempotents of $T(\rho, \preceq)$, and describe the generators of $T(\rho, \preceq)$. In Sect. 3 we describe the Green's $*$ -relations \mathcal{L}^* and \mathcal{R}^* on $T(\rho, \preceq)$. As a consequence, $T(\rho, \preceq)$ is shown to be abundant. Finally, in Sect. 4, we use the results of the previous sections to determine all automorphisms of the monoid $T(\rho, \preceq)$.

Throughout the paper, we use the following notations: let $|A|$ denote the cardinality of a set A . 1_A denote the identity function from A to itself. For a function $\alpha : A \rightarrow B$, denote the image of α by $\text{im } \alpha$. $|\text{im } \alpha|$ is said to be the *rank* of α , and we can write

$$\alpha = \begin{pmatrix} A_1 & \dots & A_k \\ a_1 & \dots & a_k \end{pmatrix},$$

where $\text{im } \alpha = \{a_1, \dots, a_k\}$, $a_i\alpha^{-1} = A_i$ ($i = 1, \dots, k$) and $\{A_1, \dots, A_k\} = A/\ker \alpha$, where $\ker \alpha$ is the kernel of α (the equivalence relation $\{(x, y) \in A \times A : x\alpha = y\alpha\}$). Denote the restriction of α to C by $\alpha|_C$ for $C \subseteq A$ and $(C)\alpha$ denotes the image set of C under α .

Let E_{n-1} denote the set of idempotents in T_n of rank $n - 1$. Every element e of E_{n-1} has the form

$$e = \begin{pmatrix} b \\ a \end{pmatrix}$$

for some $a, b \in X_n, a \neq b$, which maps b to a and x to itself for any $x \in X_n - \{b\}$ [4].

For the remainder of the paper, ρ will denote an equivalence relation on X_n and \preceq will denote a total order on the partition set X_n/ρ .

2 Units, idempotents and generators

In this section we describe the units and the idempotents of $T(\rho, \preceq)$, and determine the generators of $T(\rho, \preceq)$.

We define

$$U_\rho = \{\alpha \in T_n : (x\rho)\alpha = x\rho, x \in X_n\}.$$

Clearly, $U_\rho \subseteq T(\rho, \preceq)$. Let S_n be the symmetric group on X_n . The following lemma gives the description of the group of units of $T(\rho, \preceq)$.

Lemma 2.1 *Let $\alpha \in T_n$. Then the following statements are equivalent:*

- (1) α is a unit of $T(\rho, \preceq)$.
- (2) $\alpha \in S_n \cap T(\rho, \preceq)$.
- (3) $\alpha \in U_\rho$.

Proof Obviously (1) \Rightarrow (2).

(2) \Rightarrow (3). Suppose that $\alpha \in S_n \cap T(\rho, \preceq)$. Then $\alpha^{-1} \in S_n \cap T(\rho, \preceq)$. Given any $x \in X_n$. Take any $y \in x\rho$, we have

$$y\rho = (y\alpha\alpha^{-1})\rho \preceq (y\alpha)\rho \preceq y\rho = x\rho,$$

and so $y\alpha \in x\rho$. Thus $(x\rho)\alpha \subseteq x\rho$. To prove that $x\rho \subseteq (x\rho)\alpha$ consider $z \in x\rho$. Then we have

$$x\rho = z\rho = [(z\alpha^{-1})\alpha]\rho \preceq (z\alpha^{-1})\rho \preceq z\rho$$

and so $z\alpha^{-1} \in x\rho$ and $z = (z\alpha^{-1})\alpha$. It follows that $(x\rho)\alpha = x\rho$ and hence $\alpha \in U_\rho$.

(3) \Rightarrow (1). By the definition of U_ρ , we have that $\alpha \in T(\rho, \preceq)$ and the restriction $\alpha|_{x\rho}$ is a bijection from the ρ -class $x\rho$ onto itself, and so $\alpha \in S_n$ and $\alpha^{-1} \in T(\rho, \preceq)$. Thus α is a unit of $T(\rho, \preceq)$. \square

Next we have

Lemma 2.2 *Let $\alpha \in T(\rho, \preceq)$. Then α is an idempotent if and only if for all $t \in \text{im } \alpha$, $t\alpha = t$ and $t\rho = \min\{x\rho : x \in t\alpha^{-1}\}$.*

Proof First recall that $\alpha \in T_n$ is an idempotent if and only if, for all $t \in \text{im } \alpha$, $t \in t\alpha^{-1}$. Hence $\alpha \in T(\rho, \preceq)$ is an idempotent if and only if, for all $t \in \text{im } \alpha$,

$$t\alpha = t \quad \text{and} \quad t\rho = \min\{x\rho : x \in t\alpha^{-1}\}$$

since $t \in t\alpha^{-1}$ and $x \in t\alpha^{-1}$ implies that $t = t\alpha = x\alpha$ and $t\rho = (x\alpha)\rho \preceq x\rho$. \square

Corollary 2.3 *Let $a, b \in X_n$ and $a \neq b$. Then $a\rho \preceq b\rho$ if and only if $\binom{b}{a} \in T(\rho, \preceq)$.*

From Corollary 2.3 we introduce the following idempotent subset of $T(\rho, \preceq)$ with rank $n - 1$ (that is often used in the paper):

$$E_\rho^w = \left\{ \begin{pmatrix} b \\ a \end{pmatrix} \in E_{n-1} : a\rho \preceq b\rho \right\}.$$

Then we have

Lemma 2.4 *Let α be an element of $T(\rho, \preceq) - U_\rho$. Then α is a product of elements in $E_\rho^w \cup U_\rho$.*

Proof Let $s(\alpha)$ be the cardinality of the set $\{x \in X_n : x\alpha \neq x\}$. We will show that α can be written as a product of elements in $E_\rho^w \cup U_\rho$ by using induction on $s(\alpha)$.

Clearly, if $s(\alpha) = 1$ then $\alpha \in E_\rho^w$. We now assume that $s(\alpha) > 1$ and let

$$\alpha = \begin{pmatrix} A_1 & A_2 & \cdots & A_k \\ a_1 & a_2 & \cdots & a_k \end{pmatrix} \in T(\rho, \preceq) - U_\rho.$$

Then, for every i , $a_i\rho \preceq \min\{x\rho : x \in A_i\}$ from the definition of $T(\rho, \preceq)$. Choose $b_i \in A_i$ such that $b_i\rho = \min\{x\rho : x \in A_i\}$ for every i . Without loss of generality we may assume that

$$a_1 = b_1, \dots, a_i = b_i, \quad a_{i+1} \neq b_{i+1}, \dots, a_k \neq b_k.$$

Define

$$\beta = \begin{pmatrix} A_1 & \cdots & A_k \\ b_1 & \cdots & b_k \end{pmatrix} \quad \text{and} \quad \gamma = \begin{pmatrix} b_1 & \cdots & b_i & b_{i+1} & \cdots & b_k & Y \\ b_1 & \cdots & b_i & a_{i+1} & \cdots & a_k & y \end{pmatrix},$$

where $Y = X_n - \{b_1, \dots, b_k\}$, $y \in Y$ and $y\rho = \min\{z\rho : z \in Y\}$. Then $\beta \in T(\rho, \preceq) - U_\rho$ and $\gamma \in T(\rho, \preceq)$. Moreover, $\gamma \in U_\rho$ if and only if $k = n - 1$.

Note that $s(\alpha) = n - i > n - k = s(\beta)$ and $s(\gamma) = s(\alpha) - 1$. Thus both β and γ are products of elements in $E_\rho^w \cup U_\rho$ by inductive supposition. It follows that $\alpha = \beta\gamma$ is a product of elements in $E_\rho^w \cup U_\rho$. This completes the proof. \square

From Lemma 2.4 we immediately deduce

Proposition 2.5 *Let U_ρ and E_ρ^w be defined above. Then $E_\rho^w \cup U_\rho$ is a generating set of $T(\rho, \preceq)$.*

Similar to the proof as Lemma 2.4 we also have

Lemma 2.6 *Let α be an idempotent of $T(\rho, \preceq) - U_\rho$. Then α is a product of elements in E_ρ^w .*

3 Green's *-relations \mathcal{L}^* and \mathcal{R}^*

We recall some of the basis facts about Green's *-relations \mathcal{L}^* and \mathcal{R}^* . The relations $\mathcal{L}^*(\mathcal{R}^*)$ is defined on a semigroup S by the rule $a\mathcal{L}^*b(a\mathcal{R}^*b)$ if and only if the elements a, b of S are related by the Green's relation $\mathcal{L}(\mathcal{R})$ in some oversemigroup of S [2]. The following lemma, from [2], provides us an alternative description for $\mathcal{L}^*(\mathcal{R}^*)$.

Lemma 3.1 *Let S be a semigroup and let a, b be in S . The following conditions are equivalent:*

- (1) $a\mathcal{L}^*b(a\mathcal{R}^*b)$.
- (2) for all $s, t \in S^1$, $as = at(sa = ta)$ if and only if $bs = bt(sb = tb)$.

For $\alpha \in T_n$, $\ker \alpha = \{(x, y) \in X_n \times X_n : x\alpha = y\alpha\}$. Then we have

Proposition 3.2 *Let α, β be elements of $T(\rho, \preceq)$. Then*

- (1) $(\alpha, \beta) \in \mathcal{L}^*$ if and only if $\text{im } \alpha = \text{im } \beta$;
- (2) $(\alpha, \beta) \in \mathcal{R}^*$ if and only if $\ker \alpha = \ker \beta$.

Proof (1) Certainly if $\text{im } \alpha = \text{im } \beta$ then $(\alpha, \beta) \in \mathcal{L}(T_n)$ [5] and so $(\alpha, \beta) \in \mathcal{L}^*$.

Conversely, suppose that $(\alpha, \beta) \in \mathcal{L}^*$. Then

$$\alpha\gamma = \alpha\delta \quad \text{if and only if} \quad \beta\gamma = \beta\delta \quad (\text{for all } \gamma, \delta \in T(\rho, \preceq)).$$

Let $X_n/\rho = \{Y_1 < Y_2 < \dots < Y_t\}$. If $Y_1 = \{x\}$, for some $x \in X_n$, then, certainly $x\alpha = x = x\beta$; Otherwise, choose $y \in Y_1$ and $y \neq x$, then $\begin{pmatrix} x \\ y \end{pmatrix} \in T(\rho, \preceq)$. Hence

$$x \notin \text{im } \alpha \Leftrightarrow \alpha \cdot \begin{pmatrix} x \\ y \end{pmatrix} = \alpha \cdot 1_{X_n} \Leftrightarrow \beta \cdot \begin{pmatrix} x \\ y \end{pmatrix} = \beta \cdot 1_{X_n} \Leftrightarrow x \notin \text{im } \beta.$$

We therefore conclude that $\text{im } \alpha = \text{im } \beta$.

- (2) Again if $\ker \alpha = \ker \beta$ then $(\alpha, \beta) \in \mathcal{R}(T_n)$ [5] and so $(\alpha, \beta) \in \mathcal{R}^*$.

Conversely, if $(\alpha, \beta) \in \mathcal{R}^*$ then

$$\gamma\alpha = \delta\alpha \Leftrightarrow \gamma\beta = \delta\beta \quad (\text{for all } \gamma, \delta \in T(\rho, \preceq)).$$

For $x, y \in X_n$ with $x \neq y$, we may assume that $y\rho \preceq x\rho$. Then $\begin{pmatrix} x \\ y \end{pmatrix} \in T(\rho, \preceq)$. Hence

$$x\alpha = y\alpha \Leftrightarrow \begin{pmatrix} x \\ y \end{pmatrix} \cdot \alpha = 1_{X_n} \cdot \alpha \Leftrightarrow \begin{pmatrix} x \\ y \end{pmatrix} \cdot \beta = 1_{X_n} \cdot \beta \Leftrightarrow x\beta = y\beta.$$

We therefore conclude that $\ker \alpha = \ker \beta$. □

A semigroup S in which each \mathcal{L}^* -class and each \mathcal{R}^* -class contains an idempotent is called *abundant* [2]. We have

Corollary 3.3 For any equivalence relation ρ on X_n and a total order \preceq on the partition set X_n/ρ , the semigroup $T(\rho, \preceq)$ is abundant.

Proof For a typical \mathcal{R}^* -class R_α^* of $T(\rho, \preceq)$ with $X_n/\ker \alpha = \{A_1, A_2, \dots, A_k\}$. Choose $b_i \in A_i$ such that $b_i \rho = \min\{x_i \rho : x_i \in A_i\}$ for every $i \in \{1, \dots, k\}$. Then we have that

$$\beta = \begin{pmatrix} A_1 & \dots & A_k \\ b_1 & \dots & b_k \end{pmatrix}$$

is an idempotent of $T(\rho, \preceq)$ by Lemma 2.2, and $\alpha R^* \beta$ by Proposition 3.2.

Next, consider that a typical \mathcal{L}^* -class L_α^* of $T(\rho, \preceq)$ with $\text{im } \alpha = \{a_1, \dots, a_k\}$, where $1 \leq k \leq n$. We will use induction on k to show that L_α^* contains an idempotent of $T(\rho, \preceq)$. If $k = 1$ then, clearly α is an idempotent. Suppose now that the conclusion holds for $k - 1$. Consider that

$$\alpha = \begin{pmatrix} A_1 & \dots & A_{k-1} & A_k \\ a_1 & \dots & a_{k-1} & a_k \end{pmatrix} \in T(\rho, \preceq).$$

Without loss of generality, we may assume that

$$a_1 \rho \preceq \dots \preceq a_{k-1} \rho \preceq a_k \rho.$$

Then $\begin{pmatrix} a_k \\ a_{k-1} \end{pmatrix} \in T(\rho, \preceq)$ by Corollary 2.3, and so

$$\beta = \begin{pmatrix} A_1 & \dots & A_{k-2} & A_{k-1} \cup A_k \\ a_1 & \dots & a_{k-2} & a_{k-1} \end{pmatrix} = \alpha \begin{pmatrix} a_k \\ a_{k-1} \end{pmatrix} \in T(\rho, \preceq).$$

By induction supposition, there exists an idempotent $\varepsilon \in T(\rho, \preceq)$ such that $\beta \mathcal{L}^* \varepsilon$. Hence, by Lemma 2.2, we can let

$$\varepsilon = \begin{pmatrix} B_1 & \dots & B_{k-2} & B_{k-1} \\ a_1 & \dots & a_{k-2} & a_{k-1} \end{pmatrix}$$

with $a_i \in B_i, i = 1, \dots, k - 1$ and $a_i \rho = \min\{x_i \rho : x_i \in B_i\}$ for every $i \in \{1, \dots, k - 1\}$. Since $a_k \in X_n = B_1 \cup \dots \cup B_{k-1}$, there exist a unique B_j such that $a_k \in B_j$. Note that $a_k \neq a_j$, so $a_j \in B_j \setminus \{a_k\}$. It follows, by Lemma 2.2, that

$$\varepsilon^* = \begin{pmatrix} B_1 & \dots & B_{j-1} & B_j \setminus \{a_k\} & B_{j+1} & \dots & B_{k-1} & a_k \\ a_1 & \dots & a_{j-1} & a_j & a_{j+1} & \dots & a_{k-1} & a_k \end{pmatrix}$$

is an idempotent in $T(\rho, \preceq)$, so that $\alpha \mathcal{L}^* \varepsilon^*$ as required. \square

4 Automorphisms of $T(\rho, \preceq)$

After the preliminaries of the previous two sections, to determine every automorphism of $T(\rho, \preceq)$, we need the following lemmas.

Lemma 4.1 Let $\begin{pmatrix} b_1 \\ a_1 \end{pmatrix}, \begin{pmatrix} b_2 \\ a_2 \end{pmatrix} \in E_{n-1}$. Then $\begin{pmatrix} b_1 \\ a_1 \end{pmatrix} \begin{pmatrix} b_2 \\ a_2 \end{pmatrix} = \begin{pmatrix} b_1 \\ a_1 \end{pmatrix}$ if and only if $b_1 = b_2$.

Proof Assume that $b_1 = b_2$. Then, from $a_1 \neq b_1$ we have

$$b_1 \begin{pmatrix} b_1 \\ a_1 \end{pmatrix} \begin{pmatrix} b_1 \\ a_2 \end{pmatrix} = a_1 \begin{pmatrix} b_1 \\ a_2 \end{pmatrix} = a_1 = b_1 \begin{pmatrix} b_1 \\ a_1 \end{pmatrix},$$

and for every $x \in X_n \setminus \{b_1\}$,

$$x \begin{pmatrix} b_1 \\ a_1 \end{pmatrix} \begin{pmatrix} b_1 \\ a_2 \end{pmatrix} = x \begin{pmatrix} b_1 \\ a_2 \end{pmatrix} = x = x \begin{pmatrix} b_1 \\ a_1 \end{pmatrix}$$

and so $\begin{pmatrix} b_1 \\ a_1 \end{pmatrix} \begin{pmatrix} b_1 \\ a_2 \end{pmatrix} = \begin{pmatrix} b_1 \\ a_1 \end{pmatrix}$.

Assume now that $b_1 \neq b_2$. Then

$$b_2 \begin{pmatrix} b_1 \\ a_1 \end{pmatrix} \begin{pmatrix} b_2 \\ a_2 \end{pmatrix} = b_2 \begin{pmatrix} b_2 \\ a_2 \end{pmatrix} = a_2 \neq b_2 = b_2 \begin{pmatrix} b_1 \\ a_1 \end{pmatrix},$$

and so $\begin{pmatrix} b_1 \\ a_1 \end{pmatrix} \begin{pmatrix} b_2 \\ a_2 \end{pmatrix} \neq \begin{pmatrix} b_1 \\ a_1 \end{pmatrix}$. □

Similar argument as in Lemma 4.1, we have

Lemma 4.2 Let $\begin{pmatrix} x \\ y \end{pmatrix}, \begin{pmatrix} y \\ w \end{pmatrix} \in E_{n-1}$. Then $\begin{pmatrix} y \\ w \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$ is an idempotent if and only if $x = w$.

Lemma 4.3 Let $\begin{pmatrix} b_1 \\ a_1 \end{pmatrix}, \begin{pmatrix} b_2 \\ a_2 \end{pmatrix} \in T(\rho, \preceq)$ and $b_1 \rho \preceq b_2 \rho$. We have

- (1) If $b_1 = b_2$ then $\begin{pmatrix} b_1 \\ a_1 \end{pmatrix} \mathcal{L}^* \begin{pmatrix} b_2 \\ a_2 \end{pmatrix}$.
- (2) If $b_1 \neq b_2$ then there exists $\delta \in T(\rho, \preceq)$ such that $\begin{pmatrix} b_1 \\ a_1 \end{pmatrix} \mathcal{R}^* \delta \mathcal{L}^* \begin{pmatrix} b_2 \\ a_2 \end{pmatrix}$.

Proof Since $\text{im} \begin{pmatrix} b_1 \\ a_1 \end{pmatrix} = X_n \setminus \{b_1\} = \text{im} \begin{pmatrix} b_1 \\ a_2 \end{pmatrix}$, so (1) holds by Lemma 3.1.

For (2), note that $\begin{pmatrix} b_2 \\ b_1 \end{pmatrix} \in T(\rho, \preceq)$ by Corollary 2.3, since $b_1 \rho \preceq b_2 \rho$. It follows that $\delta = \begin{pmatrix} b_1 \\ a_1 \end{pmatrix} \begin{pmatrix} b_2 \\ b_1 \end{pmatrix} \in T(\rho, \preceq)$ with $\ker \delta = \ker \begin{pmatrix} b_1 \\ a_1 \end{pmatrix}$ and $\text{im} \delta = \text{im} \begin{pmatrix} b_2 \\ b_1 \end{pmatrix} = \text{im} \begin{pmatrix} b_2 \\ a_2 \end{pmatrix}$. Hence, by Proposition 3.2, we have $\begin{pmatrix} b_1 \\ a_1 \end{pmatrix} \mathcal{R}^* \delta \mathcal{L}^* \begin{pmatrix} b_2 \\ a_2 \end{pmatrix}$. □

Let $\text{Aut } T(\rho, \preceq)$ be the automorphism group of $T(\rho, \preceq)$. Then we have

Lemma 4.4 Let $\varphi \in \text{Aut } T(\rho, \preceq)$. Then $(E_\rho^w) \varphi = E_\rho^w$.

Proof First, we claim that

$$(E_\rho^w) \varphi \cap E_\rho^w \neq \emptyset.$$

Assume that it is not such case. Note that $(\varepsilon_i) \varphi$ is idempotent for every $\varepsilon_i \in E_\rho^w$, and so we certainly have $|\text{im}(\varepsilon_i) \varphi| \leq n - 2$. Next, given an element $\varepsilon \in E_\rho^w$ then there exists an idempotent $\omega \in T(\rho, \preceq) - \{1_{X_n}\}$ such that $(\omega) \varphi = \varepsilon$. Thus, by Lemma 2.6, there exist elements $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_k$ of E_ρ^w such that $\omega = \varepsilon_1 \varepsilon_2 \cdots \varepsilon_k$. Hence we have

$$\varepsilon = (\varepsilon_1 \varepsilon_2 \cdots \varepsilon_k) \varphi = (\varepsilon_1) \varphi (\varepsilon_2) \varphi \cdots (\varepsilon_k) \varphi.$$

It follows that $n - 1 = |\operatorname{im} \varepsilon| \leq |\operatorname{im}(\varepsilon_k)\varphi| \leq n - 2$ (since $(\varepsilon_k)\varphi \notin E_\rho^w$), a contradiction.

We have shown that $(E_\rho^w)\varphi \cap E_\rho^w \neq \emptyset$. Next, let $\varepsilon_0 = \begin{pmatrix} b_0 \\ a_0 \end{pmatrix} = \left(\begin{pmatrix} b_1 \\ a_1 \end{pmatrix}\right)\varphi \in (E_\rho^w)\varphi \cap E_\rho^w$. For any $\varepsilon = \begin{pmatrix} b \\ a \end{pmatrix} \in E_\rho^w$, we distinguish two cases as follows.

Case 1: $b = b_1$. In this case, by (1) of Lemma 4.3 we have $\varepsilon \mathcal{L}^*\begin{pmatrix} b_1 \\ a_1 \end{pmatrix}$. Further, by Lemma 3.1, we immediately deduce $(\varepsilon)\varphi \mathcal{L}^* \varepsilon_0$. Hence, by (1) of Proposition 3.2, $|\operatorname{im}(\varepsilon)\varphi| = n - 1$ and so $(\varepsilon)\varphi \in E_\rho^w$.

Case 2: $b \neq b_1$; $b\rho \leq b_1\rho$ (Similar argument for $b_1\rho \leq b\rho$). By (2) of Lemma 4.3, there exists $\delta \in T(\rho, \leq)$ such that $\varepsilon \mathcal{R}^* \delta \mathcal{L}^*\begin{pmatrix} b_1 \\ a_1 \end{pmatrix}$. Hence, by Lemma 3.1 we have $(\varepsilon)\varphi \mathcal{R}^*(\delta)\varphi \mathcal{L}^* \varepsilon_0$. That is, $\ker(\varepsilon)\varphi = \ker(\delta)\varphi$ and $\operatorname{im}(\delta)\varphi = \operatorname{im} \varepsilon_0$. It follows that

$$|\operatorname{im}(\varepsilon)\varphi| = |X_n / \ker(\varepsilon)\varphi| = |X_n / \ker(\delta)\varphi| = |\operatorname{im}(\delta)\varphi| = |\operatorname{im} \varepsilon_0| = n - 1,$$

and so $(\varepsilon)\varphi \in E_\rho^w$.

By Case 1 and Case 2 above, we have shown that $(E_\rho^w)\varphi \subseteq E_\rho^w$. By using the foregoing argument for the automorphism φ^{-1} , we have $(E_\rho^w)\varphi^{-1} \subseteq E_\rho^w$. It follows that

$$E_\rho^w = (E_\rho^w)\varphi^{-1}\varphi \subseteq (E_\rho^w)\varphi \subseteq E_\rho^w,$$

and so $(E_\rho^w)\varphi = E_\rho^w$. \square

Lemma 4.5 *Let $\varphi \in \operatorname{Aut} T(\rho, \leq)$. Then there exists $\mu_\varphi \in U_\rho$ such that $\left(\begin{pmatrix} x \\ y \end{pmatrix}\right)\varphi = \begin{pmatrix} x\mu_\varphi \\ y\mu_\varphi \end{pmatrix}$ for every $\begin{pmatrix} x \\ y \end{pmatrix} \in E_\rho^w$.*

Proof Let $X_n/\rho = \{Y_1 < Y_2 < \dots < Y_t\}$. We distinguish two cases: $|Y_1| = 1$ or $|Y_1| \geq 2$.

Case 1: $|Y_1| = 1$. Let $Y_1 = \{1\}$. For any $x \in X_n \setminus \{1\}$, clearly $\begin{pmatrix} x \\ 1 \end{pmatrix} \in E_\rho^w$ and by Lemma 4.4 we can let $\left(\begin{pmatrix} x \\ 1 \end{pmatrix}\right)\varphi = \begin{pmatrix} x' \\ 1'_x \end{pmatrix} \in E_\rho^w$. We will prove that $1 = 1'_x$. Indeed, if $1 \neq 1'_x$ then $\begin{pmatrix} 1'_x \\ 1 \end{pmatrix} \in E_\rho^w$, and by Lemma 4.4 we can let $\left(\begin{pmatrix} z \\ t \end{pmatrix}\right)\varphi = \begin{pmatrix} 1'_x \\ 1 \end{pmatrix}$, where $\begin{pmatrix} z \\ t \end{pmatrix} \in E_\rho^w$. Since $z \neq t$ and $Y_1 \leq t\rho \leq z\rho$, we have $z \neq 1$ and so

$$\begin{pmatrix} z \\ t \end{pmatrix} \begin{pmatrix} x \\ 1 \end{pmatrix} = \begin{cases} \begin{pmatrix} x \\ 1 \end{pmatrix}, & t = 1, x = z \\ \begin{pmatrix} z \\ t \end{pmatrix}, & t \neq 1, x = z \\ \begin{pmatrix} \{z, x, 1\} & k & \dots \\ 1 & k & \dots \end{pmatrix}, & t = 1, x \neq z \\ \begin{pmatrix} \{z, t\} & \{x, 1\} & k & \dots \\ t & 1 & k & \dots \end{pmatrix}, & t \neq 1, x \neq z \end{cases}$$

is an idempotent, but $\begin{pmatrix} 1'_x \\ 1 \end{pmatrix}$ is not idempotent by Lemma 4.2, a contradiction. Thus $1 = 1'_x$ and we have proved that

$$\left(\begin{pmatrix} x \\ 1 \end{pmatrix}\right)\varphi = \begin{pmatrix} x' \\ 1 \end{pmatrix} \quad \text{for any } x \in X_n \setminus \{1\}.$$

Now we define a mapping $\mu_\varphi : X_n \rightarrow X_n$ by $1\mu_\varphi = 1$ and $x\mu_\varphi = x'$ (defined the above), $x \neq 1$. Obviously, μ_φ is a bijection.

Next, we will prove that $\left(\begin{smallmatrix} x \\ y \end{smallmatrix}\right)\varphi = \begin{pmatrix} x\mu_\varphi \\ y\mu_\varphi \end{pmatrix}$ for any $\begin{pmatrix} x \\ y \end{pmatrix} \in E_\rho^w$, $x \neq 1$, $y \neq 1$. By Lemma 4.4 we can suppose that $\left(\begin{smallmatrix} x \\ y \end{smallmatrix}\right)\varphi = \begin{pmatrix} x^* \\ y^* \end{pmatrix} \in E_\rho^w$. Since

$$\begin{pmatrix} x\mu_\varphi \\ 1 \end{pmatrix} = \left(\begin{smallmatrix} x \\ 1 \end{smallmatrix}\right)\varphi = \left(\begin{smallmatrix} x \\ 1 \end{smallmatrix}\right)\begin{pmatrix} x \\ y \end{pmatrix}\varphi = \begin{pmatrix} x\mu_\varphi \\ 1 \end{pmatrix}\begin{pmatrix} x^* \\ y^* \end{pmatrix},$$

we immediately deduce that $x^* = x\mu_\varphi$ by Lemma 4.1. Moreover, from $\left(\begin{smallmatrix} y \\ 1 \end{smallmatrix}\right)\begin{pmatrix} x \\ y \end{pmatrix}\varphi = \begin{pmatrix} y\mu_\varphi \\ 1 \end{pmatrix}\begin{pmatrix} x\mu_\varphi \\ y^* \end{pmatrix}$ we see that if $y\mu_\varphi \neq y^*$ then $\begin{pmatrix} y\mu_\varphi \\ 1 \end{pmatrix}\begin{pmatrix} x\mu_\varphi \\ y^* \end{pmatrix}$ is an idempotent, and so $\begin{pmatrix} y \\ 1 \end{pmatrix}\begin{pmatrix} x \\ y \end{pmatrix}$ is idempotent (since φ is an automorphism). But this is impossible since $\begin{pmatrix} y \\ 1 \end{pmatrix}\begin{pmatrix} x \\ y \end{pmatrix}$ is not an idempotent. Hence $y\mu_\varphi = y^*$. It follows that if $y\rho \leq x\rho$ then $(y\mu_\varphi)\rho \leq (x\mu_\varphi)\rho$, and so $\mu_\varphi \in U_\rho$.

Case 2: $|Y_1| \geq 2$. In this case, for any $x \in X_n$, there exists $y \in X_n \setminus \{x\}$ such that $y\rho \leq x\rho$, and so $\begin{pmatrix} x \\ y \end{pmatrix} \in E_\rho^w$. By Lemma 4.4 we can let

$$\left(\begin{smallmatrix} x \\ y \end{smallmatrix}\right)\varphi = \begin{pmatrix} x' \\ y' \end{pmatrix} \in E_\rho^w \quad \text{and} \quad \left(\begin{smallmatrix} x \\ z \end{smallmatrix}\right)\varphi = \begin{pmatrix} x'_z \\ z' \end{pmatrix} \in E_\rho^w.$$

Note that

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \left(\begin{smallmatrix} x \\ y \end{smallmatrix}\right)\varphi = \left[\left(\begin{smallmatrix} x \\ y \end{smallmatrix}\right)\begin{pmatrix} x \\ z \end{pmatrix}\right]\varphi = \begin{pmatrix} x' \\ y' \end{pmatrix}\begin{pmatrix} x'_z \\ z' \end{pmatrix},$$

and so $x' = x'_z$ by Lemma 4.1. Hence, φ induces a map μ_φ from X_n to itself, defined by

$$\left(\begin{smallmatrix} x \\ y \end{smallmatrix}\right)\varphi = \begin{pmatrix} x\mu_\varphi \\ y' \end{pmatrix}, \quad \text{for every } \begin{pmatrix} x \\ y \end{pmatrix} \in E_\rho^w.$$

Obviously, μ_φ is surjective.

Fact 1 μ_φ is injective.

Indeed, assume that $t = x_1\mu_\varphi = x_2\mu_\varphi$, for $x_1, x_2 \in X_n$. Since $|Y_1| \geq 2$, there exist $y_1, y_2 \in X_n$ such that $\begin{pmatrix} x_1 \\ y_1 \end{pmatrix}, \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} \in E_\rho^w$. By Lemma 4.4 we let $\left(\begin{smallmatrix} x_1 \\ y_1 \end{smallmatrix}\right)\varphi = \begin{pmatrix} t \\ y'_1 \end{pmatrix}$ and $\left(\begin{smallmatrix} x_2 \\ y_2 \end{smallmatrix}\right)\varphi = \begin{pmatrix} t \\ y'_2 \end{pmatrix}$. Then

$$\left(\begin{smallmatrix} x_1 \\ y_1 \end{smallmatrix}\right)\begin{pmatrix} x_2 \\ y_2 \end{pmatrix}\varphi = \begin{pmatrix} t \\ y'_1 \end{pmatrix}\begin{pmatrix} t \\ y'_2 \end{pmatrix} = \begin{pmatrix} t \\ y'_1 \end{pmatrix} = \left(\begin{smallmatrix} x_1 \\ y_1 \end{smallmatrix}\right)\varphi.$$

As φ is a bijection it follows that $\begin{pmatrix} x_1 \\ y_1 \end{pmatrix}\begin{pmatrix} x_2 \\ y_2 \end{pmatrix} = \begin{pmatrix} x_1 \\ y_1 \end{pmatrix}$, and so $x_1 = x_2$ from Lemma 4.1.

Fact 2 If $x_1\rho \leq x_2\rho$. Then $(x_1\mu_\varphi)\rho \leq (x_2\mu_\varphi)\rho$.

In fact, assume that $(x_1\mu_\varphi)\rho \succ (x_2\mu_\varphi)\rho$. Then $\begin{pmatrix} x_2 \\ x_1 \end{pmatrix}, \begin{pmatrix} x_1\mu_\varphi \\ x_2\mu_\varphi \end{pmatrix} \in E_\rho^w$. By Lemma 4.4 we can let

$$\left(\begin{pmatrix} x_2 \\ x_1 \end{pmatrix}\right)\varphi = \begin{pmatrix} x_2\mu_\varphi \\ x_1' \end{pmatrix}, \quad \left(\begin{pmatrix} x_1 \\ z \end{pmatrix}\right)\varphi = \begin{pmatrix} x_1\mu_\varphi \\ x_2\mu_\varphi \end{pmatrix}.$$

Note that $\begin{pmatrix} x_1\mu_\varphi \\ x_2\mu_\varphi \end{pmatrix}\begin{pmatrix} x_2\mu_\varphi \\ x_1' \end{pmatrix}$ is an idempotent. Hence $\begin{pmatrix} x_1 \\ z \end{pmatrix}\begin{pmatrix} x_2 \\ x_1 \end{pmatrix}$ is also an idempotent, and so $z = x_2$ by Lemma 4.2. Further,

$$\begin{pmatrix} x_1\mu_\varphi \\ x_2\mu_\varphi \end{pmatrix} = \left(\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}\right)\varphi = \left(\begin{pmatrix} x_2 \\ x_1 \end{pmatrix}\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}\right)\varphi = \begin{pmatrix} x_2\mu_\varphi \\ x_1' \end{pmatrix}\begin{pmatrix} x_1\mu_\varphi \\ x_2\mu_\varphi \end{pmatrix}$$

is an idempotent. Hence, $x_1' = x_1\mu_\varphi$ by Lemma 4.2. Since now both $\begin{pmatrix} x_2\mu_\varphi \\ x_1\mu_\varphi \end{pmatrix}$ and $\begin{pmatrix} x_1\mu_\varphi \\ x_2\mu_\varphi \end{pmatrix}$ belong to E_ρ^w , we have $(x_1\mu_\varphi)\rho = (x_2\mu_\varphi)\rho$, which contradicts the assumption.

Fact 3 Let $\left(\begin{pmatrix} x \\ y \end{pmatrix}\right)\varphi = \begin{pmatrix} x\mu_\varphi \\ y' \end{pmatrix}$ for $\begin{pmatrix} x \\ y \end{pmatrix} \in E_\rho^w$. Then $y' = y\mu_\varphi$.

Since $\begin{pmatrix} x \\ y \end{pmatrix} \in E_\rho^w$, we have $y\rho \leq x\rho$, and so $\begin{pmatrix} x\mu_\varphi \\ y\mu_\varphi \end{pmatrix} \in E_\rho^w$ by Facts 2 and 1. By Lemma 4.4 we can let

$$\left(\begin{pmatrix} x \\ z \end{pmatrix}\right)\varphi = \begin{pmatrix} x\mu_\varphi \\ y\mu_\varphi \end{pmatrix} \quad \text{for some } z \in X_n \setminus \{x\}.$$

It is sufficient to show that $y = z$. Assume now that $y \neq z$. Then $y\rho \leq z\rho$ or $z\rho \leq y\rho$ since \leq is a totally order.

For the former, by Facts 2 and 1 we have $\begin{pmatrix} z\mu_\varphi \\ y\mu_\varphi \end{pmatrix} \in E_\rho^w$ and by Lemma 4.4 we can let $\begin{pmatrix} z\mu_\varphi \\ y\mu_\varphi \end{pmatrix} = \left(\begin{pmatrix} z \\ t \end{pmatrix}\right)\varphi$, for some $t \in X_n$. Note that $\begin{pmatrix} z\mu_\varphi \\ y\mu_\varphi \end{pmatrix}\begin{pmatrix} x\mu_\varphi \\ y\mu_\varphi \end{pmatrix}$ is idempotent. Then $\begin{pmatrix} z \\ t \end{pmatrix}\begin{pmatrix} x \\ y \end{pmatrix}$ is also idempotent, and so $t = x$ by Lemma 4.2. Hence

$$\begin{pmatrix} z\mu_\varphi \\ y\mu_\varphi \end{pmatrix} = \left(\begin{pmatrix} z \\ x \end{pmatrix}\right)\varphi = \left(\begin{pmatrix} x \\ z \end{pmatrix}\begin{pmatrix} x \\ y \end{pmatrix}\right)\varphi = \begin{pmatrix} x\mu_\varphi \\ y\mu_\varphi \end{pmatrix}\begin{pmatrix} z\mu_\varphi \\ y\mu_\varphi \end{pmatrix}.$$

It follows that $x\mu_\varphi = z\mu_\varphi$ and so $x = z$, which contradicts $x \neq z$.

For the latter, then $\begin{pmatrix} y \\ z \end{pmatrix} \in E_\rho^w$ and by Lemma 4.4 we can let $\left(\begin{pmatrix} y \\ z \end{pmatrix}\right)\varphi = \begin{pmatrix} y\mu_\varphi \\ z' \end{pmatrix}$ for some $z' \in X_n$. Note that $\begin{pmatrix} y \\ z \end{pmatrix}\begin{pmatrix} x \\ y \end{pmatrix}$ is idempotent. We have that $\begin{pmatrix} y\mu_\varphi \\ z' \end{pmatrix}\begin{pmatrix} x\mu_\varphi \\ y\mu_\varphi \end{pmatrix}$ is also idempotent, and so $z' = x\mu_\varphi$ by Lemma 4.2. Hence, from

$$\left(\begin{pmatrix} x \\ z \end{pmatrix}\begin{pmatrix} y \\ z \end{pmatrix}\right)\varphi = \begin{pmatrix} x\mu_\varphi \\ y\mu_\varphi \end{pmatrix}\begin{pmatrix} y\mu_\varphi \\ z' \end{pmatrix} = \begin{pmatrix} y\mu_\varphi \\ x\mu_\varphi \end{pmatrix} = \left(\begin{pmatrix} y \\ z \end{pmatrix}\right)\varphi$$

and φ is a bijection, we have that $\begin{pmatrix} x \\ z \end{pmatrix}\begin{pmatrix} y \\ z \end{pmatrix} = \begin{pmatrix} y \\ z \end{pmatrix}$ and so $x = y$, a contradiction.

We have proved Fact 3, and from Fact 2 we see that $\mu_\varphi \in U_\rho$, and so the proof of Lemma 4.5 is now completes. \square

Lemma 4.6 For any $\begin{pmatrix} x \\ y \end{pmatrix} \in E_\rho^w$ and for any $\mu \in U_\rho$ we have that

$$\mu^{-1}\begin{pmatrix} x \\ y \end{pmatrix}\mu = \begin{pmatrix} x\mu \\ y\mu \end{pmatrix}.$$

Proof For every $z \in X_n$ we have

$$z\mu^{-1}\binom{x}{y}\mu = \begin{cases} y\mu, & z\mu^{-1} = x \\ z, & z\mu^{-1} \neq x \end{cases} = \begin{cases} y\mu, & z = x\mu \\ z, & z \neq x\mu \end{cases} = z\binom{x\mu}{y\mu},$$

and so $\mu^{-1}\binom{x}{y}\mu = \binom{x\mu}{y\mu}$. \square

We now can state and prove the main result of the paper as follows:

Theorem 4.7 *Let S_n the symmetric group of X_n and $U_\rho = \{\mu \in S_n : (x\rho)\mu = x\rho, x \in X_n\}$. Then U_ρ is the unit group of $T(\rho, \preceq)$ and, for any $\varphi \in \text{Aut } T(\rho, \preceq)$, there exists $\mu \in U_\rho$ such that $(\alpha)\varphi = \mu^{-1}\alpha\mu$ for all $\alpha \in T(\rho, \preceq)$.*

Conversely, let $\mu \in U_\rho$. Then $\mu^{-1}\alpha\mu \in T(\rho, \preceq)$ for any $\alpha \in T(\rho, \preceq)$, and the map $\varphi : T(\rho, \preceq) \rightarrow T(\rho, \preceq)$ defined by $(\alpha)\varphi = \mu^{-1}\alpha\mu, \alpha \in T(\rho, \preceq)$ is an automorphism.

Proof Let $\varphi \in \text{Aut } T(\rho, \preceq)$. Then, first by Lemmas 4.5 and 4.6, there exists $\mu \in U_\rho$ such that

$$\left(\binom{x}{y}\right)\varphi = \binom{x\mu}{y\mu} = \mu^{-1}\binom{x}{y}\mu, \quad \text{for any } \binom{x}{y} \in E_\rho^w.$$

Secondly, we will prove that $(\alpha)\varphi = \mu^{-1}\alpha\mu$ for all $\alpha \in U_\rho$. For $\alpha \in U_\rho$, we must have $(\alpha)\varphi \in U_\rho$, and by Lemma 4.6, for any $\binom{x}{y} \in E_\rho^w$, we have

$$\begin{aligned} \binom{x\alpha\mu}{y\alpha\mu} &= \left(\binom{x\alpha}{y\alpha}\right)\varphi = \left(\alpha^{-1}\binom{x}{y}\alpha\right)\varphi \\ &= (\alpha^{-1})\varphi\left(\binom{x}{y}\right)\varphi(\alpha)\varphi = (\alpha)\varphi^{-1}\binom{x\mu}{y\mu}(\alpha)\varphi = \binom{x\mu(\alpha)\varphi}{y\mu(\alpha)\varphi}. \end{aligned}$$

Hence, $x\alpha\mu = x\mu(\alpha)\varphi$ and $y\alpha\mu = y\mu(\alpha)\varphi$. It follows that $(\alpha)\varphi = \mu^{-1}\alpha\mu$.

Finally, for any $\alpha \in T(\rho, \preceq) - E_\rho^w - U_\rho$, we have that α is a product of elements in $E_\rho^w \cup U_\rho$ by Proposition 2.5. It follows that $(\alpha)\varphi = \mu^{-1}\alpha\mu$ as required.

The converse part is obvious. \square

Remark The μ is unique in Theorem 4.7. Indeed, let μ_1, μ_2 be elements of U_ρ such that $\mu_1^{-1}\alpha\mu_1 = \mu_2^{-1}\alpha\mu_2$ for all $\alpha \in T(\rho, \preceq)$. Now, given any $x \in X_n$, we can choose an element $y \in X_n$ such that $\binom{x}{y} \in E_\rho^w$ or $\binom{y}{x} \in E_\rho^w$. For the former (Similarly for the latter), we have

$$\binom{x\mu_1}{y\mu_1} = \mu_1^{-1}\binom{x}{y}\mu_1 = \mu_2^{-1}\binom{x}{y}\mu_2 = \binom{x\mu_2}{y\mu_2}.$$

Hence $x\mu_1 = x\mu_2$, and so $\mu_1 = \mu_2$.

The map $\Psi : \text{Aut } T(\rho, \preceq) \rightarrow U_\rho$ defined by

$$(\varphi)\Psi = \mu_\varphi, \quad \varphi \in \text{Aut } T(\rho, \preceq), \quad \mu_\varphi \in U_\rho \quad \text{with } (\alpha)\varphi = \mu_\varphi^{-1}\alpha\mu_\varphi, \quad \forall \alpha \in T(\rho, \preceq)$$

is an isomorphism. We have

Corollary 4.8 $\text{Aut } T(\rho, \preceq) \cong U_\rho$.

Let PT_n be the semigroup of partial transformations on X_n . Let $S_n^- = \{\alpha \in T_n : x\alpha \leq x, \forall x \in X_n\}$ [9]. Then, by Corollary 4.8, we have

Corollary 4.9 $\text{Aut } T_n \cong S_n$; $\text{Aut } PT_n \cong S_n$ and $\text{Aut } S_n^- \cong \{1_{X_n}\}$.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

1. Araújo, J., Konieczny, J.: Automorphism groups of centralizers of idempotents. *J. Algebra* **269**, 227–239 (2003)
2. Fountain, J.B.: Abundant semigroups. *Proc. Lond. Math. Soc.* **44**(33), 103–129 (1982)
3. Ganyushkin, O., Mazorchuk, V.: Classical Finite Transformation Semigroups: an Introduction. *Algebra and Application*, vol. 9 (2009)
4. Higgins, P.: *Techniques of Semigroup Theory*. Oxford University Press, Oxford (1992)
5. Howie, J.M.: *Fundamentals of Semigroup Theory*. Oxford University Press, Oxford (1995)
6. Schreier, J.: Über Abbildungen einer Abstrakten Menge auf ihre Teilmengen. *Fundam. Math.* **28**, 261–264 (1936)
7. Sullivan, R.P.: Automorphisms of transformation semigroups. *J. Aust. Math. Soc.* **20**, 77–84 (1975)
8. Šutov, È.G.: Homomorphisms of the semigroup of all partial transformations. *Izv. Vysš. Učebn. Zaved., Mat.* **3**, 177–184 (1961)
9. Umar, A.: On the semigroups of order-decreasing finite full transformations. *Proc. R. Soc. Edinb., Sect. A, Math.* **120**, 129–142 (1992)
10. Xiuliang, Y., Haobo, Y.: Maximal half-transitive submonoids of full transformation semigroups. *Adv. Math.* **40**(5), 580–586 (2011)